

Structure of the gluon propagator at finite temperature

H. Arthur Weldon

Department of Physics, West Virginia University, Morgantown WV 26506-6315

(December 16, 1996)

Abstract

The thermal self-energy of gluons generally depends on four Lorentz-invariant functions. Only two of these occur in the hard thermal loop approximation of Braaten and Pisarski because of the abelian Ward identity $K_\mu \Pi_{\text{htl}}^{\mu\nu} = 0$. However, for the exact self-energy $K_\mu \Pi^{\mu\nu} \neq 0$. In linear gauges the Slavnov-Taylor identity is shown to require a non-linear relation among three of the Lorentz-invariant self-energy function: $(\Pi_C)^2 = (K^2 - \Pi_L)\Pi_D$. This reduces the exact gluon propagator to a simple form containing only two types of poles: one that determines the behavior of transverse electric and magnetic gluons and one that controls the longitudinally polarized electric gluons.

11.10.Wx, 12.38.Mh, 25.75.+r

I. INTRODUCTION

A. Background

The constraint imposed by local gauge invariance on the various Green functions of gauge theories are contained in the Slavnov-Taylor identities [1,2]. These identities were originally formulated in covariant gauge, which has a gauge-fixing term $\mathcal{L}_{\text{gf}} = -(\partial_\mu A^\mu)^2/2\xi$. In covariant gauge the identity satisfied by the photon propagator of QED is

$$K_\mu D'^{\mu\nu}_{\text{qed}}(K) = -\xi \frac{K^\nu}{K^2}, \quad (1.1)$$

whereas the identity satisfied by the gluon propagator of QCD is slightly weaker:

$$K_\mu K_\nu D'^{\mu\nu}(K) = -\xi. \quad (1.2)$$

At zero temperature, the abelian identity (1.1) has the same consequence as the nonabelian version (1.2) for the respective self-energy tensors. In both cases $\Pi^{\mu\nu}$ must be a linear combination of the available tensors $g^{\mu\nu}$ and $K^\mu K^\nu$. When the propagator is expressed in terms of the self-energy, application of (1.1) yields $K_\mu \Pi^{\mu\nu}_{\text{qed}} = 0$ for the abelian case and application of (1.2) yields $K_\mu \Pi^{\mu\nu} = 0$ for QCD.

At non-zero temperature the Slavnov-Taylor identities are unchanged, as shown in Appendix C. However the self-energy tensor can depend on the four-velocity u^μ of the plasma. Consequently in both QED and QCD the self-energy tensor can be a linear combination of four possible tensors: $g^{\mu\nu}$, $K^\mu K^\nu$, $u^\mu u^\nu$, and $K^\mu u^\nu + u^\mu K^\nu$. It is convenient to choose particular linear combinations of these as the standard basis set [3–5] denoted $A^{\mu\nu}$, $B^{\mu\nu}$, $C^{\mu\nu}$, $D^{\mu\nu}$ and defined in Appendix A. By parametrizing the tensor structure of the self-energy one can then calculate the form of the exact propagator and apply the Slavnov-Taylor identity. For QED the result remains $K_\mu \Pi^{\mu\nu}_{\text{qed}} = 0$, since the electromagnetic current operator satisfies $\partial_\mu j^\mu = 0$ even at non-zero temperature. Only two of the four basis tensors satisfy this. Therefore

$$\Pi^{\mu\nu}_{\text{qed}} = \Pi_T^{\text{qed}} A^{\mu\nu} + \Pi_L^{\text{qed}} B^{\mu\nu}, \quad (1.3)$$

where the subscripts T and L denote transverse and longitudinal with respect to the spatial component $\vec{k}$ of the wave-vector.

For QCD Braaten and Pisarski [6,7] discovered a consistent high-temperature approximation which behaves in the same fashion as quantum electrodynamics. In this hard thermal loop approximation the quarks are massless and only the order T^2 part of one-loop diagrams are retained. The order T^2 part of the one-loop gluon self-energy is gauge invariant and satisfies $K_\mu \Pi_{\text{htl}}^{\mu\nu} = 0$. In the standard basis $\Pi_{\text{htl}}^{\mu\nu} = \Pi_T^{\text{htl}} A^{\mu\nu} + \Pi_L^{\text{htl}} B^{\mu\nu}$, and the functions Π_T, Π_L are gauge invariant. This was discovered by explicit calculation in covariant gauges [8] and was shown by Braaten and Pisarski to be a consequence of the fact that the hard thermal loops with any number of external gluons all satisfy abelian Ward identities [6,7,9,10]. In hard thermal loop approximation the gluon propagator has the same structure as the photon thermal propagator.

If one goes beyond the hard thermal loop approximation to QCD these simplifications do not apply. Inclusion of sub-dominant powers of T , quark masses, or higher loop effects all spoil the abelian features. For example, even at one-loop order with massless quarks, the gluon self-energy has a sub-dominant term linear in T that yields $K_\mu \Pi^{\mu\nu} \neq 0$ [11]. Thus, generally

$$K_\mu \Pi^{\mu\nu} \neq 0. \quad (1.4)$$

The QCD self-energy therefore depends on all four basis tensors:

$$\Pi^{\mu\nu} = \Pi_T A^{\mu\nu} + \Pi_L B^{\mu\nu} + \Pi_C C^{\mu\nu} + \Pi_D D^{\mu\nu}, \quad (1.5)$$

where $K_\mu C^{\mu\nu} \neq 0$ and $K_\mu D^{\mu\nu} \neq 0$. In the exact self-energy the functions $\Pi_T, \Pi_L, \Pi_C, \Pi_D$ are all gauge-dependent.

B. Summary of Results

The purpose of this paper is to investigate the structure of the full gluon propagator at finite temperature that results from the self-energy (1.5). In a general linear gauge the

nonabelian Slavnov-Taylor identity (1.2) generalizes to

$$F_\mu F_\nu D'^{\mu\nu} = -\xi, \quad (1.6)$$

where F^μ is the gauge-fixing vector. The obvious way of implementing this is to insert the free propagator with gauge-fixing vector F^μ and the self-energy (1.5) into the Schwinger-Dyson equation for the full propagator $D'^{\mu\nu}$. After some tedious algebra is performed the Slavnov-Taylor constraint (1.6) can be applied.

A more direct procedure will be followed here. The quadratic part of the gluon action is the sum of the kinetic energy, the self-energy, and the gauge-fixing term. The reduced action without the gauge-fixing term will be denoted by a check:

$$\check{\Gamma}^{\mu\nu} = -K^2 g^{\mu\nu} + K^\mu K^\nu + \Pi^{\mu\nu}. \quad (1.7)$$

Sec III will show that the Slavnov-Taylor identity (1.6) can be expressed as the pair of constraints:

$$\check{\Gamma}^{\mu\nu} G_\nu = 0 \quad F^\nu G_\nu = -\xi, \quad (1.8)$$

which determine the vector G^ν . The consequences of this are simple. Viewed as a matrix, $\check{\Gamma}^\mu{}_\nu$ must have four eigenvalues. Three eigenvalues are determined dynamically by the equations of motion. The fourth eigenvalue must be zero because of local gauge invariance as expressed in (1.8). A zero eigenvalue is only possible if the determinant of $\check{\Gamma}^\mu{}_\nu$ vanishes. This immediately gives a non-linear relation on the gluon self-energy.

The condition $\text{Det}(\check{\Gamma}^\mu{}_\nu) = 0$ applies with any linear gauge-fixing condition. If we specialize to gauges that preserve the rotational invariance of the plasma rest frame, then the gauge-fixing vector F^μ should be a linear combination of K^μ and the plasma velocity u^μ . In all such cases the self-energy has the tensor structure (1.5). The requirement $\text{Det}(\check{\Gamma}^\mu{}_\nu) = 0$ leads to

$$(\Pi_C)^2 = (K^2 - \Pi_L)\Pi_D, \quad (1.9)$$

which must hold for all K^μ . This is quite remarkable since the functions Π_L, Π_C, Π_D are all gauge-dependent. In a perturbative expansion, Π_C is known to begin at order g^2 [11] and consequently Π_D must begin at order g^4 . Results equivalent to this were obtained by Gross, Pisarski, and Yaffe [3] and by Kajantie and Kapusta [4] in covariant gauge and in general linear gauges by Kobes, Kunstatter, and Mak [11].

Because of the simplification (1.9) it is rather easy to compute the full gluon propagator from the Schwinger-Dyson equation. The full propagator is constructed of four tensors but it does not have four poles. Instead there are only two: one pole associated with the two transverse polarizations contained in $A^{\mu\nu}$ that determines the behavior of transverse electric and magnetic gluons; and one pole associated with a particular combination of $B^{\mu\nu}, C^{\mu\nu}, D^{\mu\nu}$ that controls the longitudinal electric gluons.

The entire development is done in the real-time formulation of thermal QCD. The full gluon propagator has a 2×2 matrix structure $D'_{ab}{}^{\mu\nu}$, where $a, b = 1, 2$. The formulas displayed throughout this paper are for the diagonal entries of the 2×2 matrix structure. The diagonal entries satisfy Schwinger-Dyson equations that involve products of functions rather than products of matrices. For example, if k_0 is given a small positive imaginary part, i.e. $k_0 + i\epsilon$, then all formulas in the paper apply to retarded propagators and self-energies. If instead, k_0 is given an imaginary part with alternating sign, i.e. $k_0 + i\epsilon k_0$, then all formulas apply to propagators and self-energies that satisfy Feynman boundary conditions. Appendix C gives further details.

In Sec II the exact form of the thermal gluon propagator is displayed for three popular gauges: covariant, Coulomb, and temporal. The proof of these results is delayed until Sec III, which derives the general result (3.25) for the thermal gluon propagator for any linear gauge-fixing condition that preserves the rotational invariance of the plasma rest frame. Sec IV discusses the gauge invariance of the location of the transverse and longitudinal poles and shows that the two types of poles coincide at $\vec{k} = 0$. Appendix A gives explicit details about the four basis tensors. Appendix B derives the form of the gluon propagator in the hard thermal loop approximation for a general linear gauge. Appendix C derives the

Slavnov-Taylor identity for general linear gauges at finite temperature.

II. GLUON PROPAGATOR IN THREE GAUGES

The final result for the full gluon propagator in a general linear gauge is given in (3.25). Before deriving the general form it may be helpful to display the final result in familiar gauges, viz. covariant, Coulomb, and temporal. To give these results, some notation is necessary. Instead of the four-vectors K^μ and u^μ it is simpler to use K^μ and $\tilde{K}^\mu$ where

$$\tilde{K}^\mu = (K^\mu K \cdot u - u^\mu K^2)/k, \quad (2.1)$$

and $k^2 \equiv (K \cdot u)^2 - K^2 \geq 0$. This vector satisfies

$$K \cdot \tilde{K} = 0 \quad \tilde{K}^2 = -K^2. \quad (2.2)$$

It is often convenient to use the plasma rest frame $u^\mu = (1, 0, 0, 0)$ in which case $\tilde{K}^\mu = (k, \hat{k}k_0)$. In this frame the transverse tensor $A^{\mu\nu}$ has the following value

$$A^{\mu\nu}|_{\text{rest}} = \begin{pmatrix} 0 & 0 \\ 0 & -\delta^{mn} + \hat{k}^m \hat{k}^n \end{pmatrix}. \quad (2.3)$$

A second notation requires solving (1.9) for Π_C :

$$\Pi_C = \sigma \sqrt{(K^2 - \Pi_L)\Pi_D}, \quad (2.4)$$

where $\sigma = \pm 1$. In terms of this, it will be useful to employ the four-vector

$$H^\mu = K^\mu \sqrt{K^2 - \Pi_L} + \sigma \tilde{K}^\mu \sqrt{\Pi_D}, \quad (2.5)$$

which has components $H^\mu = (h_0, \hat{k} h)$.

A. Covariant gauges

In covariant gauges the free propagator is

$$D_{\text{free}}^{\mu\nu} = \left[-g^{\mu\nu} + (1 - \xi) \frac{K^\mu K^\nu}{K^2} \right] \frac{1}{K^2}. \quad (2.6)$$

or equivalently

$$D_{\text{free}}^{\mu\nu} = -\frac{A^{\mu\nu}}{K^2} + \frac{\tilde{K}^\mu \tilde{K}^\nu}{(K^2)^2} - \xi \frac{K^\mu K^\nu}{(K^2)^2}. \quad (2.7)$$

In this gauge the exact propagator (3.25) reduces to

$$D'^{\mu\nu} = \frac{-A^{\mu\nu}}{K^2 - \Pi_T} + \frac{\tilde{K}^\mu \tilde{K}^\nu}{K^2(K^2 - \Pi_L)} - \frac{\xi H^\mu H^\nu}{(K^2)^2(K^2 - \Pi_L)}. \quad (2.8)$$

At $K^2 = \Pi_T$ there is a pole in the transverse sector. At $K^2 = \Pi_L$ there is a pole which appears to have a complicated tensor structure. However, as $K^2 \rightarrow \Pi_L$ the auxiliary vector $H^\mu \rightarrow \sigma\sqrt{\Pi_D}\tilde{K}^\mu$. Consequently at this pole

$$(K^2 - \Pi_L)D'^{\mu\nu}\big|_{\text{pole}} \rightarrow [1 - \xi \frac{\Pi_D}{K^2}] \frac{\tilde{K}^\mu \tilde{K}^\nu}{K^2}, \quad (2.9)$$

and the non-transverse function Π_D only modifies the residue of the pole.

B. Coulomb gauges

A second example is provided by generalized Coulomb gauges with gauge-fixing term $\mathcal{L}_{\text{gf}} = -(\nabla \cdot \mathbf{A})^2/2\xi$. The free gluon propagator is

$$D_{\text{free}}^{\mu\nu} = -\frac{A^{\mu\nu}}{K^2} + \frac{1}{k^2} \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} - \xi \frac{K^\mu K^\nu}{k^4}. \quad (2.10)$$

In this case the D^{00} propagator is instantaneous and only the transverse potentials have time-dependence. The true Coulomb gauge $\nabla \cdot \mathbf{A} = 0$ corresponds to $\xi = 0$. The full propagator that results from (3.25) is

$$D'^{\mu\nu} = \frac{-A^{\mu\nu}}{K^2 - \Pi_T} + \frac{K^2}{h^2} \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} - \xi \frac{H^\mu H^\nu}{h^2 k^2}, \quad (2.11)$$

where h is the spatial component of H^μ :

$$h = k\sqrt{K^2 - \Pi_L} + \sigma k_0\sqrt{\Pi_D}. \quad (2.12)$$

As $h \rightarrow 0$, $H^\mu \rightarrow (h^0, 0)$ and $h^0 \rightarrow -\sigma\sqrt{\Pi_D}K^2/k$. Consequently this pole also has a simple tensor structure:

$$h^2 D'^{\mu\nu}\big|_{\text{pole}} \rightarrow K^2[1 - \xi\Pi_D \frac{K^2}{k^4}] \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}. \quad (2.13)$$

C. Temporal gauges

A third type of example is the generalized temporal gauge propagator. The gauge-fixing term is $\mathcal{L}_{\text{gf}} = -(A^0)^2/2\xi$, where ξ has dimensions of length squared. The free propagator is

$$D_{\text{free}}^{\mu\nu} = -\frac{A^{\mu\nu}}{K^2} + \frac{1}{k_0^2} \begin{pmatrix} 0 & 0 \\ 0 & \hat{k}^m \hat{k}^n \end{pmatrix} - \xi \frac{K^\mu K^\nu}{k_0^2}. \quad (2.14)$$

The exact propagator (3.25) reduces in this gauge to

$$D'^{\mu\nu} = -\frac{A^{\mu\nu}}{K^2 - \Pi_T} + \frac{K^2}{h_0^2} \begin{pmatrix} 0 & 0 \\ 0 & \hat{k}^m \hat{k}^n \end{pmatrix} - \xi \frac{H^\mu H^\nu}{h_0^2}, \quad (2.15)$$

where h_0 is the time component of H^μ :

$$h_0 = k_0 \sqrt{K^2 - \Pi_L} + \sigma k \sqrt{\Pi_D}. \quad (2.16)$$

At momenta for which $h_0 \rightarrow 0$, $H^\mu \rightarrow (0, \hat{k}h)$ and $h \rightarrow \sigma \sqrt{\Pi_D} K^2/k_0$ so that the tensor structure of the residue is given by:

$$h_0^2 D'^{\mu\nu} \rightarrow K^2 [1 - \xi \Pi_D \frac{K^2}{k_0^2}] \begin{pmatrix} 0 & 0 \\ 0 & \hat{k}^m \hat{k}^n \end{pmatrix} \quad (2.17)$$

As before Π_D modifies the numerical value of the residue but not the spin structure.

III. PROOF FOR GENERAL LINEAR GAUGES

A. Gauge fixing condition

The investigation pertains to any linear gauge-fixing condition that is linear and that does not break rotational invariance in the plasma rest frame. With a gauge-fixing term $\mathcal{L}_{\text{gf}} = -(\mathcal{F}_\mu A^\mu)^2/2\xi$ the quadratic part of the momentum-space action is

$$\frac{1}{2} \int \frac{d^4 K}{(2\pi)^4} A_\mu(K) \Gamma_{\text{free}}^{\mu\nu} A_\nu(-K), \quad (3.1)$$

where

$$\Gamma_{\text{free}}^{\mu\nu} = -K^2 g^{\mu\nu} + K^\mu K^\nu - \frac{F^\mu F^\nu}{\xi}. \quad (3.2)$$

The most general F^μ that does not spoil the rotational invariance is a linear combination of two vectors:

$$F^\mu = f_1 K^\mu + f_2 \tilde{K}^\mu, \quad (3.3)$$

where f_1 and f_2 are functions of k_0 and k . In the plasma rest frame $K^\mu = (k_0, \vec{k})$ and $\tilde{K}^\mu = (k, k_0 \hat{k})$. It will be helpful to also introduce the vector

$$\tilde{F}^\mu = f_1 \tilde{K}^\mu + f_2 K^\mu, \quad (3.4)$$

with the property

$$F \cdot \tilde{F} = 0. \quad (3.5)$$

For the covariant, Coulomb, and temporal gauges the vectors (3.3) and (3.4) are as follows:

	F^μ	$\tilde{F}^\mu$	f_1	f_2
Covariant	$(k_0, \vec{k})$	$(k, k_0 \hat{k})$	1	0
Coulomb	$(0, \vec{k})$	$(k, 0)$	$-k^2/K^2$	$k_0 k/K^2$
Temporal	$(1, 0)$	$(0, \hat{k})$	k_0/K^2	$-k/K^2$

(3.6)

The free propagator is the inverse of (3.2):

$$D_{\text{free}}^{\mu\nu} = -\frac{A^{\mu\nu}}{K^2} + \frac{\tilde{F}^\mu \tilde{F}^\nu}{(f_1 K^2)^2} - \xi \frac{K^\mu K^\nu}{(f_1 K^2)^2}. \quad (3.7)$$

B. Exact gluon propagator

For the full gluon propagator it is necessary to add the self-energy to (3.2) to obtain the quadratic part of the full effective action:

$$\Gamma^{\mu\nu} = -K^2 g^{\mu\nu} + K^\mu K^\nu + \Pi^{\mu\nu} - \frac{F^\mu F^\nu}{\xi}. \quad (3.8)$$

The full gluon propagator, denoted $D'_{\nu\alpha}$ is the inverse

$$\Gamma^{\mu\nu} D'_{\nu\alpha} = \delta^\mu_\alpha. \quad (3.9)$$

The full propagator must satisfy the nonabelian Slavnov-Taylor identity

$$F^\nu F^\alpha D'_{\nu\alpha} = -\xi, \quad (3.10)$$

which is the generalization of (1.2) to general linear gauges. Appendix C shows that this identity must hold even at finite temperature. To determine how (3.10) directly constrains the proper self-energy, it is useful to define a four-vector G_ν by

$$G_\nu \equiv D'_{\nu\alpha} F^\alpha. \quad (3.11)$$

Then (3.9) implies that

$$\Gamma^{\mu\nu} G_\nu = F^\mu, \quad (3.12)$$

and (3.10) implies

$$F^\nu G_\nu = -\xi. \quad (3.13)$$

Consequently (3.12) can be written

$$\left(\Gamma^{\mu\nu} + \frac{F^\mu F^\nu}{\xi} \right) G_\nu = 0. \quad (3.14)$$

The tensor in parentheses will be denoted by a check:

$$\begin{aligned} \check{\Gamma}^{\mu\nu} &\equiv \Gamma^{\mu\nu} + F^\mu F^\nu / \xi \\ &= -K^2 g^{\mu\nu} + K^\mu K^\nu + \Pi^{\mu\nu}. \end{aligned} \quad (3.15)$$

In terms of the tensor decomposition (1.5) the explicit form is

$$\begin{aligned} \check{\Gamma}^\mu{}_\nu &= (\Pi_T - K^2) A^\mu{}_\nu + (\Pi_L - K^2) B^\mu{}_\nu \\ &\quad + \Pi_C C^\mu{}_\nu + \Pi_D D^\mu{}_\nu. \end{aligned} \quad (3.16)$$

Although $\check{\Gamma}$ has the gauge-fixing term removed it is still gauge-dependent because of the self-energy functions. It must satisfy

$$\check{\Gamma}^\mu_\nu G^\nu = 0, \quad (3.17)$$

where G^ν is an unknown four-vector satisfying (3.13). The first step in implementing this is to require that $\check{\Gamma}$ have determinant zero. Since $\check{\Gamma}$ is a 4×4 matrix, it must have four eigenvectors. Two of the eigenvectors are the two transverse polarization vectors ϵ^ν which satisfy $K \cdot \epsilon = 0$ and $u \cdot \epsilon = 0$. The eigenvalue for each of these is the same: $\check{\Gamma}^\mu_\nu \epsilon^\nu = (\Pi_T - K^2)\epsilon^\mu$. The remaining two eigenvectors are each linear combinations of K^ν and $\tilde{K}^\nu$. The properties of the basis tensors A, B, C, D give

$$\begin{aligned} \check{\Gamma}^\mu_\nu K^\nu &= \Pi_C \tilde{K}^\mu + \Pi_D K^\mu \\ \check{\Gamma}^\mu_\nu \tilde{K}^\nu &= (\Pi_L - K^2) \tilde{K}^\mu - \Pi_C K^\mu, \end{aligned} \quad (3.18)$$

which yields a quadratic equation for the remaining two eigenvalues. The product of all four eigenvalues is

$$\text{Det}(\check{\Gamma}^\mu_\nu) = (\Pi_T - K^2)^2 [(\Pi_L - K^2)\Pi_D + (\Pi_C)^2]. \quad (3.19)$$

The Slavnov-Taylor identity forces the quantity in square brackets to vanish for all K^μ :

$$(\Pi_L - K^2)\Pi_D + (\Pi_C)^2 = 0. \quad (3.20)$$

It seems most natural to solve this for Π_C :

$$\Pi_C = \sigma \sqrt{(K^2 - \Pi_L)\Pi_D}, \quad (3.21)$$

where $\sigma = \pm 1$. When this condition is substituted into (3.16), the particular linear combination of the tensors B, C, D can be written in terms of a single four-vector $\tilde{H}^\mu$ as follows:

$$\begin{aligned} \check{\Gamma}^{\mu\nu} &= (\Pi_T - K^2)A^{\mu\nu} + \frac{\tilde{H}^\mu \tilde{H}^\nu}{K^2} \\ \tilde{H}^\nu &\equiv \tilde{K}^\nu \sqrt{K^2 - \Pi_L} + \sigma K^\nu \sqrt{\Pi_D}. \end{aligned} \quad (3.22)$$

The null vector of $\check{\Gamma}^\mu_\nu$ is the vector

$$H^\nu = K^\nu \sqrt{K^2 - \Pi_L} + \sigma \tilde{K}^\nu \sqrt{\Pi_D}. \quad (3.23)$$

It satisfies $\check{\Gamma}^\mu_\nu H^\nu = 0$ because $\tilde{H} \cdot H = 0$. (The null vector G^ν normalized by (3.13) is $G^\nu = -\xi H^\nu / F \cdot H$.)

Now that the Slavnov-Taylor constraint has been satisfied, it is straightforward to obtain the full propagator. First add back the gauge-fixing term to $\check{\Gamma}$:

$$\Gamma^{\mu\nu} = (\Pi_T - K^2)A^{\mu\nu} + \frac{\tilde{H}^\mu \tilde{H}^\nu}{K^2} - \frac{F^\mu F^\nu}{\xi}. \quad (3.24)$$

It is easy to invert this using $F \cdot \tilde{F} = H \cdot \tilde{H} = 0$ and $\tilde{F} \cdot \tilde{H} = -F \cdot H$ to obtain the full propagator:

$$D'^{\mu\nu} = -\frac{A^{\mu\nu}}{K^2 - \Pi_T} + K^2 \frac{\tilde{F}^\mu \tilde{F}^\nu}{(F \cdot H)^2} - \xi \frac{H^\mu H^\nu}{(F \cdot H)^2}. \quad (3.25)$$

Naturally it satisfies the Slavnov-Taylor identity (3.10). Sec II has displayed examples of this result in covariant, Coulomb, and temporal gauges.

IV. DISCUSSION

A. Gauge invariance of poles

The tensor structure of the gluon propagator (3.25) is explicitly dependent upon ξ and upon the gauge-fixing parameters f_1 and f_2 contained in F^μ and $\tilde{F}^\mu$. Moreover the self-energy $\Pi^{\mu\nu}$ is dependent upon these same gauge parameters since they occur in the free propagator (3.7). Thus all four functions Π_T , Π_L , Π_C , and Π_D are gauge-dependent.

Kobes, Kunstatter, and Rebhan proved a remarkable theorem [12]. If the exact propagator is expressed as a linear combination of the four basis tensors $A^{\mu\nu}$, $B^{\mu\nu}$, $C^{\mu\nu}$, $D^{\mu\nu}$ then the location of the poles in the $A^{\mu\nu}$ and $B^{\mu\nu}$ part of the propagator are gauge-fixing independent. The transverse projection of (3.25) is

$$A_{\mu\nu} D'^{\mu\nu} = -\frac{2}{K^2 - \Pi_T}. \quad (4.1)$$

The complex momentum that satisfy $K^2 = \Pi_T$ are gauge invariant by the KKR theorem. The $B_{\mu\nu}$ projection of (3.25) is

$$B_{\mu\nu}D'^{\mu\nu} = (\xi\Pi_D - f_1^2 K^2) \frac{K^2}{(F \cdot H)^2}. \quad (4.2)$$

The denominator is

$$F \cdot H = K^2 \left[f_1 \sqrt{K^2 - \Pi_L} - \sigma f_2 \sqrt{\Pi_D} \right]. \quad (4.3)$$

This vanishes at complex K that satisfy

$$K^2 = \Pi_L + \left(\frac{f_2}{f_1} \right)^2 \Pi_D. \quad (4.4)$$

Using (3.21) this can also be written in terms of Π_C as $K^2 = \Pi_L + \Pi_C(f_2/f_1)$. Despite the explicit gauge dependent coefficients f_1 and f_2 , the theorem of Kobes, Kunstatter, and Rebhan [12] shows that the solution of this complex dispersion relation is gauge invariant. At momenta which satisfy (4.4), $H^\mu \rightarrow \tilde{F}^\mu \sqrt{K^2 - \Pi_L}/f_1$. Therefore the tensor structure of the propagator at the pole is simple:

$$(F \cdot H)^2 D'^{\mu\nu} \Big|_{\text{pole}} \rightarrow [K^2 - \frac{\xi}{f_1^2} (K^2 - \Pi_L)] \tilde{F}^\mu \tilde{F}^\nu. \quad (4.5)$$

B. Zero Three-Momentum

The pole in the transverse sector occurs at $K^2 = \Pi_T$. The pole in the longitudinal sector occurs at (4.4). It is easy to show that at $\vec{k} = 0$ these poles coincide and that they have the same residue. This comes about because the proper self-energy $\Pi^{\mu\nu}(k_0, \vec{k})$ has the property that when $|\vec{k}| = 0$, it does not depend upon $\hat{k}$. The proof of this is quite simple. The self-energy is computed by integrations over fermion and boson propagators. Let P^α denote a generic loop momenta to be integrated and K^α denote the external four-momentum of the gluon. The fermion and boson propagators to be integrated over have arguments of the form $P^\alpha + c_j K^\alpha$, where c_j depends upon where the propagator occurs in the diagram. At

$|\vec{k}| = 0$, the arguments of the propagators are $(p_0 + c_j k_0, \vec{p})$ and all information about the direction $\hat{k}$ disappears.

To apply this result, begin with the general self-energy

$$\Pi^{\mu\nu}(k_0, \vec{k}) = \Pi_T A^{\mu\nu} + \Pi_L B^{\mu\nu} + \Pi_C C^{\mu\nu} + \Pi_D D^{\mu\nu}. \quad (4.6)$$

According to Appendix A the $|\vec{k}| = 0$ limit of the four basis tensors gives for the spatial components of the self-energy

$$\begin{aligned} \Pi^{mn}(k_0, 0) &= (-\delta^{mn} + \hat{k}^m \hat{k}^n) \Pi_T(k_0, 0) \\ &\quad - \hat{k}^m \hat{k}^n \Pi_L(k_0, 0). \end{aligned} \quad (4.7)$$

Since this cannot depend upon $\hat{k}$

$$\Pi_T(k_0, 0) = \Pi_L(k_0, 0). \quad (4.8)$$

Similarly $\Pi^{0n}(k_0, 0) = \hat{k}^n \Pi_C(k_0, 0)$ and since this cannot depend on $\hat{k}$ it is necessary that $\Pi_C(k_0, 0) = 0$. Because of relation (3.21) this automatically means that $\Pi_D(k_0, 0) = 0$. Thus the self-energy tensor is

$$\Pi^{\mu\nu}(k_0, 0) = \begin{pmatrix} 0 & 0 \\ 0 & -\delta^{mn} \Pi_T(k_0, 0) \end{pmatrix}. \quad (4.9)$$

The full gluon propagator is quite simple in this limit. For example, in covariant gauge the propagator is

$$D^{\mu\nu}(k_0, 0) = \begin{pmatrix} -\xi/k_0^2 & 0 \\ 0 & \delta^{mn}/[k_0^2 - \Pi_T(k_0, 0)] \end{pmatrix}. \quad (4.10)$$

It is also simple in the temporal gauge. In the Coulomb gauge the limit $|\vec{k}| \rightarrow 0$ does not exist for either the free or the full propagator.

ACKNOWLEDGMENTS

This work was supported in part by National Science Foundation grants PHY-9213734 and PHY-9630149.

APPENDIX A: TENSOR BASIS

There is a standard set of four tensors [3–5] constructed out of $g^{\mu\nu}$, K^μ , and $\tilde{K}^\mu = (K \cdot u K^\mu - K^2 u^\mu)/k$ that will be used throughout:

$$\begin{aligned} A^{\mu\nu} &= g^{\mu\nu} - B^{\mu\nu} - D^{\mu\nu} \\ B^{\mu\nu} &= -\frac{\tilde{K}^\mu \tilde{K}^\nu}{K^2} \\ C^{\mu\nu} &= \frac{K^\mu \tilde{K}^\nu + \tilde{K}^\mu K^\nu}{K^2} \\ D^{\mu\nu} &= \frac{K^\mu K^\nu}{K^2} \end{aligned} \tag{A1}$$

The squares of these tensors are

$$A^2 = A, \quad B^2 = B, \quad C^2 = -B - D, \quad D^2 = D \tag{A2}$$

Note that requiring A , B , and D to be idempotent fixes their sign and magnitude, but there is no such universal choice for C . The twelve mixed products are all simple:

$$\begin{aligned} AB &= AC = AD = BD = 0 \\ BA &= CA = DA = DB = 0 \\ (BC)^{\mu\nu} &= (CD)^{\mu\nu} = \tilde{K}^\mu K^\nu / K^2 \\ (CB)^{\mu\nu} &= (DC)^{\mu\nu} = K^\mu \tilde{K}^\nu / K^2 \end{aligned} \tag{A3}$$

It is often convenient to work in the rest frame of the plasma, where $u^\mu = (1, 0, 0, 0)$ and the vector $\tilde{K}^\mu$ takes the simple form $(k, \hat{k} k_0)$. In this frame the four basis tensors have the following values:

$$\begin{aligned} A^{\mu\nu} \Big|_{\text{rest}} &= \begin{pmatrix} 0 & 0 \\ 0 & -\delta^{mn} + \hat{k}^m \hat{k}^n \end{pmatrix} \\ B^{\mu\nu} \Big|_{\text{rest}} &= \frac{1}{K^2} \begin{pmatrix} -k^2 & -k_0 k^n \\ -k^m k_0 & -k_0^2 \hat{k}^m \hat{k}^n \end{pmatrix} \end{aligned}$$

$$\begin{aligned}
C^{\mu\nu}|_{\text{rest}} &= \frac{1}{K^2} \begin{pmatrix} 2k_0 k & (k_0^2 + k^2) \hat{k}^n \\ \hat{k}^m (k_0^2 + k^2) & 2k_0 k \hat{k}^m \hat{k}^n \end{pmatrix} \\
D^{\mu\nu}|_{\text{rest}} &= \frac{1}{K^2} \begin{pmatrix} k_0^2 & k_0 k^n \\ k^m k_0 & k^m k^n \end{pmatrix}
\end{aligned} \tag{A4}$$

APPENDIX B: HARD THERMAL LOOP APPROXIMATION

Although the structure of the gluon propagator in the hard thermal loop approximation is well known in standard gauges, it is useful to obtain it in a general linear gauge using the method employed in Sec III. Braaten and Pisarski showed that the gluon propagator in the hard thermal loop approximation obeys abelian Ward identities [6,7,9,10]. In a general linear gauge the abelian Ward identity is

$$D_{\nu\alpha}^{\text{htl}} F^\alpha = -\frac{\xi}{F \cdot K} K_\nu, \tag{B1}$$

as shown in Appendix C. This constraint can be rewritten by multiplying (B1) by $\Gamma_{\text{htl}}^{\mu\nu}$ to obtain

$$F^\mu = -\frac{\xi}{F \cdot K} \Gamma_{\text{htl}}^{\mu\nu} K_\nu. \tag{B2}$$

This is equivalent to

$$\left(\Gamma_{\text{htl}}^{\mu\nu} + \frac{F^\mu F^\nu}{\xi} \right) K_\nu = 0. \tag{B3}$$

As before, denote the tensor in parenthesis by a check: $\check{\Gamma}_{\text{htl}}^{\mu\nu} \equiv \Gamma_{\text{htl}}^{\mu\nu} + F^\mu F^\nu / \xi$. In terms of the tensor decomposition (1.5)

$$\begin{aligned}
\check{\Gamma}_\nu^{\mu \text{htl}} &= (\Pi_T^{\text{htl}} - K^2) A^\mu{}_\nu + (\Pi_L^{\text{htl}} - K^2) B^\mu{}_\nu \\
&+ \Pi_C^{\text{htl}} C^\mu{}_\nu + \Pi_D^{\text{htl}} D^\mu{}_\nu.
\end{aligned} \tag{B4}$$

Condition (B3) requires $\check{\Gamma}_\nu^{\mu \text{htl}} K^\nu = 0$. From the tensor definitions in Appendix A, $A^\mu{}_\nu K^\nu = 0$ and $B^\mu{}_\nu K^\nu = 0$ so that $\check{\Gamma}_\nu^{\mu \text{htl}} K^\nu = \Pi_C^{\text{htl}} \tilde{K}^\mu + \Pi_D^{\text{htl}} K^\mu$. Because the vectors $\tilde{K}^\mu$ and K^μ are

linearly independent, the only solution is $\Pi_C^{\text{htl}} = 0, \Pi_D^{\text{htl}} = 0$. To obtain the propagator in the hard thermal loop approximation, we must add the gauge-fixing term to (B4):

$$\Gamma_{\text{htl}}^{\mu\nu} = (\Pi_T^{\text{htl}} - K^2)A^{\mu\nu} + (\Pi_L^{\text{htl}} - K^2)B^{\mu\nu} - \frac{F^\mu F^\nu}{\xi}. \quad (\text{B5})$$

The inverse of this is the hard thermal loop propagator. Using $F \cdot \tilde{F} = K \cdot \tilde{K} = 0$ and $F \cdot K = -\tilde{F} \cdot \tilde{K} = f_1 K^2$ gives the result

$$D_{\text{htl}}^{\prime\mu\nu} = \frac{-A^{\mu\nu}}{K^2 - \Pi_T^{\text{htl}}} + \frac{\tilde{F}^\mu \tilde{F}^\nu}{(f_1)^2 K^2 (K^2 - \Pi_L^{\text{htl}})} - \frac{\xi K^\mu K^\nu}{(f_1 K^2)^2}. \quad (\text{B6})$$

APPENDIX C: SLAVNOV-TAYLOR IDENTITY IN GENERAL LINEAR GAUGES

This Appendix will derive the Slavnov-Taylor identity for the gluon propagator at finite temperature. The generating functional for the real-time Green functions of finite-temperature QCD is [5]

$$Z(J) = \int D\Omega \exp(iS_J), \quad (\text{C1})$$

where

$$S_J = \int_C d^4x (\mathcal{L} - \frac{1}{2\xi} (\mathcal{F}_\nu A^\nu)^2 + J_\nu A^\nu). \quad (\text{C2})$$

The label C denotes a contour in the complex time plane that runs from an initial time t_0 and ends at $t_0 - i\beta$ along a path that will be specified later. The gauge-fixing vector $\mathcal{F}_\nu$ determines the Fadeev-Popov matrix

$$\mathcal{M}(x, y) = -\mathcal{F}_{x\nu} D_x^\nu \delta^4(x - y) \quad (\text{C3})$$

and the invariant integration measure

$$D\Omega = D\psi D\bar{\psi} DA \det(\mathcal{M}). \quad (\text{C4})$$

The generating functional is invariant under a change of the integration variables A^μ . If the change of variables is a gauge transformation $A^\mu \rightarrow A^\mu + D^\mu \Lambda$ then the measure $D\Omega$ and

the quark-gluon lagrangian $\mathcal{L}$ will not change. The only change comes from the gauge-fixing and source terms. Thus

$$0 = \int D\Omega \delta S_J \exp(iS_J) \quad (\text{C5})$$

where

$$\delta S_J = \int_C d^4x \left(-\frac{(\mathcal{F}_\mu A^\mu) \mathcal{F}_\nu D^\nu \Lambda}{\xi} + J_\nu D^\nu \Lambda \right). \quad (\text{C6})$$

It is convenient to choose Λ as

$$\Lambda(x) = \int_C d^4y [\mathcal{F}_\mu D^\mu]^{-1}(x, y) \lambda(y) \quad (\text{C7})$$

for an arbitrary function $\lambda(y)$ so that

$$\frac{\delta S_J}{\delta \lambda(y)} = -\frac{1}{\xi} \mathcal{F}_\nu A^\nu(y) + \int_C d^4x J_\nu(x) D_x^\nu [\mathcal{F} \cdot D]^{-1}(x, y) \quad (\text{C8})$$

The invariance (C5) becomes

$$\frac{\mathcal{F}_\nu(y)}{Z(0)} \frac{i\delta Z}{\delta J_\nu(y)} = \int_C d^4x J_\nu(x) G^\nu(x, y) \quad (\text{C9})$$

$$G^\nu(x, y) \equiv -\xi \int D\Omega D_x^\nu [\mathcal{F} \cdot D]^{-1}(x, y) \exp(iS_J) / Z(0). \quad (\text{C10})$$

This has the property that

$$\mathcal{F}_\mu(x) G^\mu(x, y) \Big|_{J=0} = -\xi \delta_C^4(x - y). \quad (\text{C11})$$

Now apply $\delta/\delta J_\mu(x)$ to (C9)

$$\frac{\mathcal{F}_\nu(y)}{Z(0)} \frac{i\delta^2 Z}{\delta J_\mu(x) \delta J_\nu(y)} = G^\mu(x, y) + \int_C d^4z J_\alpha(z) \frac{\delta G^\alpha(z, y)}{\delta J_\mu(x)} \quad (\text{C12})$$

and set $J = 0$:

$$\mathcal{F}_\nu(y) D'^{\mu\nu}(x, y) = G^\mu(x, y) \Big|_{J=0}. \quad (\text{C13})$$

In QCD the covariant derivative in (C10) depends on the vector potential. The Slavnov-Taylor identity follows from applying $\mathcal{F}_{x\mu}$ to (C13):

$$\mathcal{F}_\mu(x)\mathcal{F}_\nu(y)D^{\mu\nu}(x,y) = -\xi \delta_C^4(x-y). \quad (\text{C14})$$

To convert this to momentum space it is necessary to know where the time components of x and y lie on the contour C in the complex time plane. For field theory in Minkowski space the contour C is taken as the union of four parts [5] that depends on a parameter σ with $0 \leq \sigma \leq \beta$: C_1 runs from $-t_0$ to t_0 , C_3 runs from t_0 to $t_0 - i\sigma$, C_2 runs from $t_0 - i\sigma$ to $-t_0 - i\sigma$, and C_4 runs from $-t_0 - i\sigma$ to $-t_0 - i\beta$. As $t_0 \rightarrow \infty$ the contributions of C_3 and C_4 can be neglected. The physical field $A_1^\mu(x) \equiv A^\mu(x)$ and the auxiliary field $A_2^\mu(x) \equiv A^\mu(t - i\sigma, \vec{x})$ lie on C_1 and C_2 , respectively. Since the time components of x and y may be taken along C_1 or C_2 independently, the propagator becomes a 2×2 matrix in the internal space. The contour-ordered propagator contains four time-ordered parts. The explicit form of (C14) is

$$\mathcal{F}_\mu(x)\mathcal{F}_\nu(y)D_{ij}^{\prime\mu\nu}(x,y) = -\xi \delta^4(x-y)(\tau_3)_{ij} \quad (\text{C15})$$

where now x and y have real time components. In momentum space this becomes

$$F_\mu F_\nu D_{ij}^{\prime\mu\nu}(K) = -\xi(\tau_3)_{ij}. \quad (\text{C16})$$

For covariant gauges $F^\mu = K^\mu$, for Coulomb gauges $F^\mu = (0, \vec{k})$, and for temporal gauge $F^\mu = (1, 0, 0, 0)$.

Feynman Basis: The 2×2 matrix structure of the time-ordered thermal propagator may be diagonalized in terms of functions which have Feynman-like analytic structure (analytic in the upper half-plane of complex $(k_0)^2$) as follows [5,13]:

$$D_{ij}^{\prime\mu\nu}(K) = U_{i\ell} \begin{pmatrix} D_F^{\prime\mu\nu}(K) & 0 \\ 0 & -D_F^{\prime\mu\nu*}(K) \end{pmatrix}_{\ell m} V_{mj}, \quad (\text{C17})$$

where U and V are nonsingular matrices that satisfy $U\tau_3V = \tau_3$. The matrix equation (C16) therefore reduces to the single condition

$$F_\mu F_\nu D_F^{\prime\mu\nu}(K) = -\xi. \quad (\text{C18})$$

Condition (C16) imposes no additional constraint on the off-diagonal parts of the propagator because $D_{12}^{\prime\mu\nu} \propto \text{Im} D_{11}^{\prime\mu\nu}$. Therefore $F_\mu F_\nu D_{12}^{\prime\mu\nu} \propto F_\mu F_\nu \text{Im} D_{11}^{\prime\mu\nu} \propto \text{Im} \xi = 0$.

Retarded Basis: The time-ordered propagator may also be diagonalized in terms of retarded propagators (analytic in the upper half-plane of complex k_0) and advanced propagators (analytic in the lower half-plane of complex k_0) [14]:

$$D'_{ij}{}^{\mu\nu}(K) = U_{i\ell} \begin{pmatrix} D'_R{}^{\mu\nu}(K) & 0 \\ 0 & -D'_A{}^{\mu\nu}(K) \end{pmatrix}_{\ell m} V_{mj} \quad (\text{C19})$$

where U and V are different from those in (C17) but still satisfy $U\tau_3 V = \tau_3$. As before the matrix equation (C16) reduces to a single condition on the retarded function

$$F_\mu F_\nu D'_R{}^{\mu\nu}(K) = -\xi. \quad (\text{C20})$$

Abelian Ward Identity: For an abelian theory the covariant derivative in (C10) is independent of the vector potential so that (C13) becomes

$$\mathcal{F}_\nu(y) D'^{\mu\nu}(x, y) = -\xi \frac{\partial}{\partial x_\nu} [\mathcal{F} \cdot \partial]^{-1}(x, y). \quad (\text{C21})$$

In momentum space this is

$$F_\nu D'_{ij}{}^{\mu\nu}(K) = -\xi \frac{K^\mu}{F \cdot K} (\tau_3)_{ij}, \quad (\text{C22})$$

which may be diagonalized in either the Feynman or the retarded basis.

REFERENCES

- [1] A.A. Slavnov, Theor. Math. Phys. **10**, 153 (1972).
- [2] J.C. Taylor, Nucl. Phys. **B33**, 436 (1971).
- [3] D.J. Gross, R.D. Pisarski, and L.G. Yaffe, Rev. Mod. Phys. **53**, 43 (1981).
- [4] K. Kajantie and J. Kapusta, Ann. Phys. (N.Y.) **160**, 477 (1985).
- [5] N.P. Landsman and Ch.G. van Weert, Phys. Rep. **145**, 141 (1987).
- [6] E. Braaten and R.D. Pisarski, Nucl. Phys. **B337**, 569 (1990) and **B339**, 310 (1990).
- [7] E. Braaten and R.D. Pisarski, Phys. Rev. **D45**, 1827 (1992).
- [8] O.K. Kalishnikov and V.V. Klimov, Phys. Lett. **B95**, 234 (1980); H. A. Weldon, Phys. Rev. D **26**, 1394 (1982).
- [9] J. Frenkel and J.C. Taylor, Nucl. Phys. **B 334**, 199 (1990).
- [10] J.C. Taylor and S.M.H. Wong, Nucl. Phys. **B346**, 115 (1990); J. Frenkel and J.C. Taylor, Nucl. Phys. **B 374**, 156 (1992).
- [11] R. Kobes, G. Kunstatter, and K.W. Mak, Z. Phys. C **45**, 129 (1989).
- [12] R. Kobes, G. Kunstatter, and A. Rebhan, Phys. Rev. Lett. **64**, 2992 (1990); Nucl. Phys. **B335**, 1 (1991).
- [13] M.A. van Eijck, R. Kobes, and Ch. G. van Weert, Phys. Rev. D **50**, 4097 (1994).
- [14] P. Aurenche and T. Becherrawy, Nucl. Phys. **B379**, 259 (1992); M.A. van Eijck and Ch. G. van Weert, Phys. Lett. **B278**, 305 (1992).